



International Journal of Agricultural and  
Environmental Research  
**FREE AND OPEN ACCESS**  
Available online at [www.ijaer.com](http://www.ijaer.com)  
ISSN 2414-8245 (Online)  
ISSN 2518-6116 (Print)



## INFLUENCE OF POTASSIUM LEVELS ON GLADIOLUS CORMS PRODUCTION IN OPEN FIELD AND INTERCROPPING CONDITIONS

FAROOQ<sup>1</sup>, ASHFAQ ALAM<sup>2</sup>, NOOR UL AMIN<sup>2</sup>, WASIM BILAL<sup>1</sup>, MOHAMMAD ABDUL RAUF<sup>1</sup>,  
KHALIL UR RAHMAN<sup>1</sup>, ROSHAN ALI<sup>1</sup>, MUHAMMAD IBRAHIM<sup>1</sup>, JAVED RAHMAN<sup>1</sup>,  
MUHAMMAD SHAHID<sup>1</sup> AND NADIA BOSTAN<sup>1</sup>.

<sup>1</sup>Agricultural Research Institute, Mingora, Swat, Khyber Pakhtunkhwa

<sup>2</sup>The University of Agriculture, Peshawar, Khyber Pakhtunkhwa

Correspondence author E-mail: [javedmiandam@gmail.com](mailto:javedmiandam@gmail.com).

### Abstract

The influence of potassium levels on gladiolus corm (*Gladiolus grandiflorus*) production under two growing conditions were studied at Agriculture Research Institute, Mingora Swat. The two growing conditions (open field and intercropping) and potassium levels (0, 30, 60 and 90) were studied during the experiment. The experiment was planned in randomized complete block design having split plot arrangement with three replications. Light intensity and soil humidity were also recorded in both conditions. Data was recorded on various growth attributes, which showed that both growing conditions and potassium levels had significantly affected most of the attributes. The maximum sprouting percentage (93.7%), diameter of corms (5.3 cm), corms weight (69.9 g), number of cormels plant<sup>-1</sup> (19.0) and survival percentage (88.8%) were recorded in open field, while maximum plant height (120.2 cm) was observed in intercropping. Moreover, maximum number of leaves plant<sup>-1</sup> (6.5), plant height (120.4 cm), diameter of corms (4.3 cm), corms weight (47.9 g), number of cormels plant<sup>-1</sup> (16.9) and survival percentage (88.8%) were recorded in plots applied with 90 kg potassium ha<sup>-1</sup>. Hence, it was concluded that gladiolus could be grown in open field condition with application of potassium @ 90 kg ha<sup>-1</sup> for better growth and production of corms under the agro-climatic conditions of Swat valley.

**Key words:** *Gladiolus*, Growing conditions and Potassium levels.

### INTRODUCTION

Gladiolus (*Gladiolus grandiflorus*) belongs to family *iridaceae*. It is indigenous to South Africa, having around one hundred and fifty known species (Negi et al., 1982). Commonly, it is known as "Sword Lily", an ornamental cormelous flower (Memon, 2009). It is cultivated on a wide range of tropical to subtropical regions of the world. Commercially gladiolus is grown from its corms for the production of both spikes and corms. In each season daughter corms are produced by the mother corms (Memon et al. 2009). Gladiolus produce cormels once the flower spikes initiate and are produced at full bloom. The size of the produced cormels increased with the downward flow of photosynthates (Hartmann et al., 1981).

Light plays a vital role in plant growth, as it is an important factor of photosynthesis and without it plant cannot prepare their own food. Plants grown under the shed receive light in less amount, which may resultantly affect their growth (Imaizumi and Kay, 2006). Moreover, plants under the shed may have less

photosynthetic activities and chlorophyll content, which may also result in less dry matter production (Rao and Mitra, 1998).

The quality of gladiolus flower is an important factor considered in the market. Hence, quality and production of the flowers is influenced by some factors such as corm size, fertilizer management and time of plantation (Arora and Khanna, 1990). Vegetative growth of a plant depends on the quantity and availability of essential nutrients in the soil. Potassium as an important nutrient, directly influences the growth, development and quality of gladiolus plant. Most soils are deficient in available potassium next to nitrogen and phosphorus, (Salisbury and Ross, 1992). Potassium has a major role in regulation of stomata and also maintains water balance in plant tissues. Furthermore, it assists the growth of meristematic tissue, involves in some enzymatic activities, responsible for metabolism of nitrogen, synthesis of proteins, used as catalyst for other reaction, accelerating metabolism of carbohydrate and works in translocation process (Bhandal and Malik, 1988).

Potassium as a catalyst plays a vital role in number of biosynthetic processes of photosynthesis. It also help plant to fight against many diseases. The quality of flower is affected in the absence of potassium even though if the plant is provided with adequate dose of nitrogenous fertilizer (Mukhopadhyay, 1995).

Pierson (1990) in a study recorded high tillers, leaves and biomass in plants grown under direct sunlight as compared to the plants under shade. Moreover, Zhao et al. (2012) recorded highest morphological attributes such as plant height and

## MATERIALS AND METHOD

The influence of potassium levels on gladiolus corm production in open field and intercropping was evaluated at Agriculture Research Institute Mingora, Swat during the spring season 2014. An exotic gladiolus variety “Amsterdam” was used in the experiment. The corms were planted in two different growing conditions in March (2014) at open field and

**Experimental design:** The experiment was conducted in the Horticulture Research area, Agricultural Research Institute (ARI), Mingora Swat. The experiment was laid out in randomized complete block design (RCBD) having split plot arrangement, replicated three times. The experiment was composed of 2 factors such as growing conditions and potassium levels. Two growing conditions (open field and intercropping) were allotted to main plots while four levels of potassium (control, 30, 60 and 90 Kg ha<sup>-1</sup>) were kept in the sub plots. Randomly selected composite soil samples were taken from the site before the experiment and were analyzed in soil and fertility lab, ARI, Mingora Swat for various physico-chemical

number of leaves in plants under direct sun light than the plants under shade. Ullah et al. (2016) found significant effect of potassium levels on the vegetative and reproductive attributes of gladiolus cultivars. The quality and yield of cotton plant were enhanced by application of potassium (Aneela et al., 2003).

Keeping in view the importance of light and potassium fertilization, this study was designed to evaluate the influence of growing condition and potassium levels on corm production of gladiolus.

in peach orchard as intercropped while, various potassium levels were applied as another factor. Nitrogen and phosphorous were applied at constant dose (120+90 kg ha<sup>-1</sup>) to all the treatments including control. Whereas, Urea was used as a source of nitrogen, SSP was taken as a source of phosphorus and SOP was used as a source of potassium. The field was thoroughly prepared and all the standard cultural practices were applied to the experimental field. The distance between row to row and plant to plant were kept 60 and 10 cm, respectively.

properties (Table 1). Various attributes were studied for all the replicated treatments and their mean values were finally recorded.

**Procedure for data recording:** All the data was recorded by adopting standard procedures. Soil moisture and light intensity were observed twice a day (Morning and afternoon) throughout the experiment and the data was recorded by Luster leaf rapitest soil moisture meter. Furthermore, sprouting percentage was determined from the number of sprouted cormels and total cormels planted by the following formula.

$$\text{Sprouting percentage} = \frac{\text{No. of cormels sprouted}}{\text{Total No. of cormels planted}} \times 100$$

**Table-1: Physico-chemical analysis of soil**

Parameters	Unit	Range	Mean	STD
pH		5.4-7.8	6.9	0.6
Lime	%	2.6-19.3	6.3	3.4
Organic Matter	%	1.2-3.4	2.2	0.6
P	mgkg <sup>-1</sup>	Mehlic No.3 (6.6-39.8) AB-DTPA (40-153)	19.1	8.7
		Mehlic No.3 (6.6-39.8) AB-DTPA (40-153)	6.3	2.9
K	mgkg <sup>-1</sup>	3.2-65.6	149.0	56.1
		26.4-78.4	88.6	29.4
Sand	%	6.4-26.8	30.7	13.8
Silt	%		59.2	12.9
Clay	%		9.9	3.1

The leaf area was measured in cm<sup>2</sup> by leaf area meter. Moreover, plant height was measured in cm using inch tape from soil surface to the apex of the plant. Diameter of corm was recorded in cm using digital vernier

caliper. Corms weight was determined in g using electronic balance while, number of cormels plant<sup>-1</sup> were observed by counting the total number of cormels produced in randomly selected plants of each

treatment. All the recorded data of each parameter was taken by randomly selecting 6 plants in each replication of each treatment.

**Statistical procedure:** The collected data was subjected to analysis of variance techniques, computed by Statistics 8.1 software and means of significant findings were further calculated according to LSD test ( $P \leq 0.05$ ) (Steel and Torri, 1997).

**Soil Physico-chemical properties:** The mean pH range, lime content, organic matter, P, K and different soil separates along with textural classes are given in Table-1.

## RESULTS AND DISCUSSION

**Light intensity and Soil Moisture:** The light intensity (166.94-foot candle) was significantly higher

**Table-2: Soil moisture and light intensity of growing conditions.**

Month	Light Intensity		Soil Moisture	
	Open field	Intercropping	Open field	Intercropping
March	107.00	53.69	222	298
April	103.44	51.3772	230	316
May	104.53	51.8108	223	294
June	174.70	51.8108	234	300
July	212.23	94.89	219	274
August	203.63	91.45	210	280
September	182.40	82.95	207	256
October	247.62	109.04	201	257
<b>Mean</b>	<b>166.94</b>	<b>76.89</b>	<b>218.25</b>	<b>284.38</b>

**Sprouting percentage:** Sprouting percentage was considerably affected only by growing conditions. The sprouting percentage (93.7) of open field condition was relatively higher than the plants sown in intercropping condition (91.7) (Table-3). The maximum sprouting percentage recorded in open field might be due to the difference in growing conditions because gladiolus is a light loving plant. Light helps in production of optimum photosynthates and dry matter that are necessary for growth and development. These findings are supported by Hangarter, (1997) and Winslow, (1999). Sprouting percentage was decreased by unfavorable environmental condition such as low light (Hong et al., 1989). The rate and proportion of germination was higher in seeds sown under light than in darkness (Colbach, 2002). The germination of seeds either sown fresh or buried in soil for several months is speed up by light (Froud-Williams, 1981; Lonchamp et al., 1984).

**Leaf area (cm<sup>2</sup>):** The growing conditions and potassium levels had significant effect on leaf area.

in open field than the light intensity (76.89-foot candle) recorded in intercropping condition. Whereas, soil moisture was maximum (284.38%) in intercropping as compared to soil moisture (218.25%) of open field (Table-1). The maximum light intensity in the open field and minimum in intercropping may be due to the fact that trees intercept different levels of light and light availability is completely affected by the shading effect of the trees around (Canham et al., 1994)). Similarly, shades create inactive photo synthetically light fall on soil surface (Fetcher et al., 1983; Turton & Duff, 1992). Soil moisture was higher in shaded area than in the open field and the same was also found by Tanga et al. (2014). The reason for this change was reduction in evapotranspiration and increase in relative humidity in the shaded area due to low light intensity (Kessler, 1992).

However, their interaction was non-significant (Table-3). The maximum leaf area (71.4 cm<sup>2</sup>) was found in

plants of intercropping, while the minimum leaf area (61.3 cm<sup>2</sup>) was observed at open field plants. The highest leaf area (69.3 cm<sup>2</sup>) was recorded in plants treated with 90 kg potassium ha<sup>-1</sup> in contrast to the lowest leaf area (62.6 cm<sup>2</sup>) of control treatment.

Leaf length increase with every consecutive increase in shade levels, as 50% shade produced significantly longer leaves than control (Kumar et al., 2013). Ramachandrudu and Thangam, (2009) found increase in leaf length and width in plants kept in shade than in open field which justify the current findings. The leaf size can be influenced by intensity of the light (Jeong et al., 2009; Vendrame et al., 2004). The plants under shade produce dry matter content in less amount as most of the food is allocated for production of shoots and have maximum leaf area (Grime, 1979). Pierson et al. (1990) found that plant exposed to full sunlight had lowest leaf area while plant grown under the shade of 60 and 90% had the highest leaf area.

Potassium plays a vital role in opening and closing of stomata for photosynthesis, which helps in cell growth and division, thus increases the area of leaf (Sober et al., 1981). The average leaf area was increased with the increasing levels of potassium application (Azizi, 1998; Asif et al., 2007).

**Plant height (cm):** Plant height was influenced by growing conditions and potassium levels. Whereas, the interaction was non-significant (Table-3). The plants grown under condition of intercropping had the maximum plant height (120.2 cm) as compared to the lowest plant height (113.5 cm) in open field. The application of 90 kg potassium ha<sup>-1</sup> resulted with maximum plant height of 120.4 cm, followed by (118.4 cm) plant treated with 30 kg potassium ha<sup>-1</sup> while the minimum plant height was noted in control treatment. The plants grown under the shade tends to gain height in search of light. In a study the plants kept under shade

of 25% were recorded to have 70% taller than the plants under direct sunlight (Zervoudakis et al., 2012). Moreover, the plants under intensity of low light were noticed maximum height (Wang et al., 2009). Another study showed that plants were taller in shade than those grown under open field conditions (Kumar et al., 2013).

The increase in plant height recorded in current study might be due to the application of potassium because potassium is an important nutrient, which helps in translocation of carbohydrates and regulates various physiological and biochemical activities that are responsible for better growth of the plants (Marschner, 1995). The plant height was increased with the application of potassium (Hussain et al., 2014). Moreover, Uddin et al. (2013) in a study on rice found that plant height was enhanced by application of potassium.

**Table-3: Sprouting percentage, leaf area and plant height as influenced by growing conditions and potassium application.**

Growing conditions	Sprouting percentage	Leaf area (cm <sup>2</sup> )	Plant height (cm)
Open field	93.7 a	61.3 b	113.5 b
Intercropping	91.7 b	71.4 a	120.2 a
LSD	1.1	8.21	4.56
<b>Potassium (Kg ha<sup>-1</sup>)</b>			
0	91.9 a	62.6 c	112.6 c
30	92.4 a	66.2 b	118.4 ab
60	92.8 a	67.3 ab	116.0 bc
90	93.8 a	69.3 a	120.4 a
LSD	NS	2.98	3.50
Interaction	NS	NS	NS

**Diameter of Corms (mm):** The two growing condition and application of potassium levels significantly varied the diameter of corms while the interaction was found non-significant (Table-4). The maximum diameter of corm (5.3 cm) was observed in plants grown in open field condition while the minimum corm diameter (2.7 cm) was noted in plant under intercropping condition. The highest level of potassium level (90 kg ha<sup>-1</sup>) produced highest corm diameter (4.3 cm), followed by corm diameter (4.0 cm) produced by application of 60 kg ha<sup>-1</sup>. Whereas, the least diameter of corm was found in control treatment. Light intensity is one of the important factors influencing the rate of photosynthesis. Photosynthesis helps in many plant processes such as metabolism and provides energy that fuels these processes, which are directly related with the plant's ability to grow. Therefore, the maximum diameter observed in plants grown in open field of the current study might be due the more photosynthetic activity. As corms are the food storage organs of gladiolus, so

the maximum food produced during the photosynthesis process in the presence of sun light have been stored in the corms that might have increased diameter of the corms.

Potassium plays an important role in corm size because potassium has a great role in the availability of other nutrients, water retention and makes roots strong. Therefore, the maximum diameter of corms was obtained in plants treated with 90 kg potassium ha<sup>-1</sup> as compare to control. Similar findings were also observed by (Baloch et al., 1991; Sing and Verma, 2001; Bybordi and Makakouti, 2003) that bulb yield was increased by increasing the level of potassium application. The bulb diameter as well as bulb quality were significantly affected by additional dose of potassium application (Desuki et al., 2006).

**Corm weight (g):** The results showed that corms weight was significantly affected by growing conditions and potassium levels; the interaction among

growing conditions and potassium levels was significant (Table-4).

The maximum corms weight (69.9 g) weight were recorded in plants of open field, whereas the minimum corms weight (19.9 g) were recorded in intercropping condition. The maximum corms weight (47.9 g) was taken with application of 90 kg potassium ha<sup>-1</sup>, followed by (45.7 g) with 60 kg potassium ha<sup>-1</sup>. The control treatment was recorded with minimum corms weight (42.2 g). The interaction between growing conditions and potassium levels showed that the greatest corms weight (72.3 g) was recorded in plants treated with 90 kg potassium ha<sup>-1</sup> and planted at open field while the minimum corms weight (16.5 g) was observed in control treatment and planted in intercropping. The maximum corms weight in open field than intercropping might be the influence of light. As light is the major source of photosynthesis process, which help in improving growth and development of Gladiolus (Young, 2003). Ramachandrudu and Thangam, (2009) also mentioned that the corms weight was maximum in open field than the weight of corms in shade, which justifies the findings of present research.

Increase dose of potassium significantly improved the corms weight (Das, 1998). The bulb weight as well as bulb quality were significantly affected by potassium application (Desuki et al., 2006).

**Number of cormels plant<sup>-1</sup>:** Number of cormels plant<sup>-1</sup> was significantly influenced by growing conditions and potassium levels, whereas the interaction between growing conditions and potassium levels was non-significant (Table-4). In case of growing conditions, the maximum number of cormels plant<sup>-1</sup> (19.0) were recorded in plants of open field while the minimum number of cormels plant<sup>-1</sup> (7.3) were observed at intercropping. The maximum number of cormels plant<sup>-1</sup> (16.9) was counted in plants applied with 90 kg potassium ha<sup>-1</sup>, followed by (13.0) with 30 kg potassium ha<sup>-1</sup>. The minimum number of cormels plant<sup>-1</sup> (10.6) were taken in control. Ramachandrudu and Thangam (2009) stated that gladiolus plant prefer sun light for better growth and development. Moreover, the results of his findings showed that maximum cormels plant<sup>-1</sup> were taken in plants grown in open field while minimum were recorded in plants under the shade. Potassium helps in the high number of cormels production. An increase in yield of cormels was observed with application of fertilizers, particularly with potassium (Misra and Singh, 1998). In addition, cormels plant<sup>-1</sup> was increased with the increase of potassium concentration (Singh et al., 1997; Barma et al., 1998). Number of researchers such as (Niwuzhang and Linx, 1997; Jiang et al., 1998; Sharma et al., 1992; Bybordi and Makakouti, 2003) reported that bulb yield was increased by increasing the level of potassium application.

**Table-4: Diameter of corm, corms weight and number of corms plant<sup>-1</sup> as influenced by growing conditions and potassium application.**

Growing conditions	Diameter of corm (cm)	Corms weight (g)	Number of cormels plant <sup>-1</sup>
Open field	5.3 a	69.9 a	19.0 a
Intercropping	2.7 b	19.9 b	7.3 b
LSD	0.43	5.87	4.28
<b>Potassium (Kg ha<sup>-1</sup>)</b>			
0	3.8 c	42.2 d	10.6 b
30	3.9 bc	43.8 c	13.0 b
60	4.0 b	45.7 b	12.1 b
90	4.3 a	47.9 a	16.9 a
LSD	0.19	0.80	2.93
Interaction	NS	1.13	NS

## REFERENCES

Aneela, S., A. Muhammad and M.E. Akhtar. 2003. Effect of potash on boll characteristics and seed cotton yield in newly developed highly resistant cotton varieties. Pak. J. Biol. Sci., 6: 813.

Arora, J.S. and K. Khanna. 1990. Studies on corm production in gladiolus as affected by different corm sizes. Indian J. Hort. 47: 442-446.

Asif, M., Amanullah and M. Anwar. 2007. Phenology, Leaf Area and Yield of Spring Maize (Cv. Azam) as affected by levels and timings of potassium application. World Applied Sci. J., 2 (4): 299-303.

Azizi, M. 1998. The effect of different irrigation regimes and potassium fertilizer on agronomic, physiological, and biochemical properties of soybean. PhD thesis in Agriculture, Mashad University., 143.

Baloch, M.A., A.F. Baloch, G. Baloch, A.H. Ansari and S.M. Qayyum, 1991. Growth and yield response of onion to different nitrogen and potassium fertilizer combination levels. Sarhad J. Agric., 7: 63-66.

Barma, G., S. Chanda and N. Roychowdhury. 1998. Production of corms and cormels of gladiolus through application of nitrogen, phosphorus and potassium. Hort. J., 11(2): 87-92.

Bhandal, I.S. and C.P. Malik. 1988. Potassium estimation, uptake, and its role in the physiology and metabolism of

- flowering plants. Intern. Review of cytology., 110: 205-254.
- Bybordi and M.J. Malakouti. 2003. The effect of various rates of potassium, zinc and copper on the yield and quality of onion under saline conditions in two major onion growing region so feast Azarbayjan. Agric. Sci. and Technol., 17: 43-52.
- Canham, C.D., A.C. Finzi, S.W. Pacala and D.H. Burbank. 1994. Causes and consequences of resource heterogeneity in forests: interspecific variation in light transmission by canopy trees. Canadian J. Forestry Res., 24: 337-349.
- Colbach, N., B. Chauvel, C. Durr and G. Richard. 2002. Effect of environmental conditions on *Alopecurus myosuroides* germination. Effect of temperature and light. Weed Res., 42: 210-221.
- Das, T.K. 1998. Corm and cormel production in gladiolus as affected by spike removal and K application. Indian J. Hort., 55(4): 327-331.
- EL-Desuki, M., M.M. Abdel-Mouty and A.H. Ali. 2006. Response of onion plants to additional dose of potassium application. J. Applied Sci., 2(9): 592-597.
- Fetcher, N., B.R. Strain and S.F. Oberbauer. 1983. Effects of light regime on the growth, leaf morphology, and water relations of seedlings of two species of tropical trees. Oecologia., 58: 314-319.
- Froud-Williams, R.J. 1981. Germination behaviour of Bromus species and *Alopecurus myosuroides*. In: Proceedings 1981 Association of Applied Biologists Conference: Grass Weeds in Cereals in the United Kingdom, Reading, UK., 31-40.
- Grime, J.P. 1979. Plant strategies and vegetation processes. John Wiley, New York.
- Hangarter, R.P. 1997. Gravity light and plant form. Plant Cell Environ., 20: 796-800.
- Hartmann, H.T., W.J. Flocker and A.M. Kofrank. 1981. Ornamental grown from bulbs, corms, tubers and rhizomes. In "Plant Sci. Growth, Dev. & Utilization of Cultiv. Plants". pp. 429-453.
- Hong, Y.P., D.H. Goo and K.Y. Huh. 1989. Studies on corm formation in gladiolus gandavensis. The effect of planting date of cormels on corm production, dormancy and flowering of the corm in the next generation. Rep. Rural Dev. Adm. Hort. Manual., 31(4): 54-59.
- Hussain, M.A., N.U. Amin, G. Ayub and M. Sajid. 2014. Response of Tuberose (*Polianthes tuberosa*) to potassium and planting depth. J. Biology, Agric. and Health care., 4(11): 106-110.
- Pierson, E.A., R.N. Mack and R.A. Black. 1990. The effect of shading on photosynthesis, growth and regrowth following defoliation for *Bromus tectorum*. Oecologia Springer-Verlag., 84: 534-543.
- Ramachandrudu, K. and M. Thangam. 2009. Production Technology of Gladiolus in Goa. Technical Bulletin No.20, ICAR Research Complex for Goa, Ela, Old Goa-403 402, Goa, India. Pp: 1-31.
- Rao, L.J., and B.N. Mitra. 1998. Growth and yield of peanut as influenced by degree and duration of shading. J. Agron Crop Sci. 160: 260-265.
- Imaizumi, T. and S.A. Kay. 2006. Photoperiodic control of flowering: not only by coincidence. Trends Plant Sci., 11:550-558.
- Jeong, K.Y., C.C. Pasian, M. McMahan and D. Tay. 2009. Growth of six Begonia species under shading. Open Hort. J., 2: 22-28.
- Jiang, Q., H. Zhang, A.I. Guanghai, Q.S. Jiang, H.L. Zhang and A.I. Gh, 1998. Application of potassium fertilizer in onion production. China-vegetables. No. 4.
- Young, J.B. and B.G.S. Jr. 2003. Field grown cut flower production in southern Louisiana. Louisiana State University, A. S. University of Louisiana at Lafayette.
- Kessler, J.J. 1992. The influence of Karite (*vitellaria paradox*) and Nere (*Parkia biglobosa*) trees on sorghum production in Burkina Faso. Agro forest. Syst., 17: 97-134.
- Kumar, R., S. Sharma, K. Ramesh and B. Singh. 2013. Effects of shade regimes and planting geometry on growth, yield and quality of the natural sweetener plant stevia (*Stevia rebaudiana Bertoni*) in north-western Himalaya. Archives of Agron. and Soil Sci., 59(7): 963-97.
- Lonchamp, J.P., R. Chadoeuf and G. Barralis. 1984. Evolution de la capacite de germination des semences de mauvaises herbes dans le sol. Agronomie., 4: 671-682.
- Marschner, H. 1995. Mineral Nutrition of Higher Plants. 2<sup>nd</sup> edition. Academic Press, London, UK. Pp. 889.
- Memon, N., M. Qasim, M.J. Jaskani, R. Ahmad and I. Ahmad. 2009. Enhancement of corm and cormel production in gladiolus (*Gladiolus* spp.). New Zealand J. Crop and Hortic. Sci., 37: 319-325.
- Misra, R.L. and B. Singh. 1998. Gladiolus. In: Commercial Flowers. (Eds) Bose, T. K. and L. P. Yadav. Naya Prokash, Calcutta, India., 266-353.
- Mukhopadhyay, A. 1995. Gladiolus. Publication and Information Division. Indian Council of Agric. Res., Krishi Anusandhan Bhavan, New Delhi. pp. 1-83.
- Memon, N., M. Qasim, M.J. Jaskani, R. Ahmad and I. Ahmad. 2009. Enhancement of corm and cormel production in gladiolus. 37 (4):319-325.
- Memon, N. N., M. Qasim, M. J. Jaskani, R. Ahmad and R. Anwar. 2009. Effect of various corm sizes on the vegetative, floral and corm yield attributes of gladiolus. Pak. J. Agri. Sci., 46:13-19.
- Negi, S.S., T.V.R.S. Sharma, S.P.S. Raghava and V.R. Shrinivasan. 1982. Variability studies in gladiolus. Indian J. Hort., 39: 269-272.
- Niwuzhang, Z.Y. and A. Linx. 1997. The effect of K source on yield and quality of some vegetable crops. Acta Agric. Zhe Jiangensis., 9: 143-148.
- Salisbury, F.B. and C. W. Ross. 1992. Mineral nutrition in Plant Physiology. 4th Ed. 116-135. Wadsworth Pub. Co. Belmont, California.
- Sharma, R.P. 1992. Effect of planting materials, nitrogen and potash on bulb yield of rainy season onion (*Allium cepa* L.). Indian, J. Agron., 37: 268-284.
- Sing, S.P. and A.B. Verma. 2001. Response of onion (*Allium cepa* L.) to potassium application. Indian J. Agron., 46: 182-185.
- Singh, K.P., N. Ramachandran and S. Uma. 1997. Growth, flowering, corm yield and corm-rot incidence as affected by level and frequency of potassium application

- in gladiolus (*Gladiolus grandiflorus*). Ind. J. Agric. Sci. 67(9): 404-406.
- Sober, M. and S.M. Zanaty. 1981. Effectiveness of inoculation with silicate bacteria in relation to the potassium content of plant using the intensive cropping technique. Res J. Agric. and Bio. Sci., 59: 280-287.
- Steel, R.G.D., J.H. Torri and D.A. Dicky. 1997. Principles and procedures of statistics. A biometric approach. 3rd ed. McGraw Hill Book Co. Inc., New York, USA.
- Tanga, A.A., T.F. Erenso and B. Lemma. 2014. The influence of Karite (*vitellaria paradox*) and Nere *Parkia biglobosa* tress on sorghum production in Burkina Faso. J. soil sci. and environ. management., 5(5): 62-71.
- Turton, S.M. and G. A. Duff. 1992. Light environments and floristic composition across an open forest-rainforest boundary in northeastern Queensland. Australian J. Ecology., 17: 415-423.
- Uddin, S., M.A.R. Sarkar and M.M. Rahman. 2013. Effect of nitrogen and potassium on yield of dry direct seeded rice cv. Nerica 1 in Ausseason. International J. Agronomy and Plant Production., 4: 69-75.
- Ullah, G., G. Ayub, M. Ilyas, I.H. Khan, M.A. Hussain, F. Naeem, Luqman and M.I. Khan. 2016. Effect of various potassium levels on growth and development of different gladiolus cultivars. Int. J. Biosci., 8(5): 45-49.
- Vendrame, W., K.K. Moore, and T.K. Broschat. 2004. Interaction of light intensity and controlled- release fertilization rate on growth and flowering of two New Guinea impatiens cultivars. Hort. Technology., 14: 491-495.
- Wang, Y., Q. Guo and M. Jin. 2009. Effects of light intensity on growth and photosynthetic characteristics of *Chrysanthemum morifolium*. Zhongguo Zhongyao Zazhi., 34: 1633-1635.
- Winslow, R.B. and E. Huala. 1999. Blue-light photoreceptors in higher plants. Annu. Rev. Cell Dev. Biol., 15: 33-62.
- Zervoudakis, G., G. Salahas, G. Kaspiris and E. Konstantopoulou. 2012. Influence of light intensity on growth and physiological characteristics of common sage (*Salvia officinalis* L.). Braz. arch. biol. technol. vol.55 no.1 Curitiba Jan./Feb. 2012.
- Zhao, D., Z. Hao and J. Tao. 2012. Effects of shade on plant growth and flower quality in the herbaceous peony (*Paeonia lactiflora* Pall.). Plant Physiol. and Biochem., 61: 187-196.